



## PRS® Stage 1 & 2 Instructions

### DIRECTIONS FOR USE:

1. Make sure the filter is clean and equipment is working properly.
2. Using a time-metered pump or by hand, add PRS Stage 1 across the pool surface at a rate of 4 oz per 20,000 gallons of pool water. For best dispersion, add product in front of the pool return lines.
3. Six hours after adding PRS Stage 1, add PRS Stage 2 across the pool surface at a rate of 4 oz per 20,000 gallons of pool water, using a time-metered pump or by hand. For best dispersion, add product in front of the pool return lines.

**Note:** Do not add both PRS Stage 1 and PRS Stage 2 at the same time.

### FREQUENCY:

**High bather load-** Use product daily. **Medium bather load-** Use product two to four times per week. **Low bather load** Use product weekly.

### IN CASE OF ACCIDENTAL FECAL RELEASE (AFR):

1. Evacuate the pool.
2. Remove all visible fecal material possible.
3. Follow local health department guidelines for proper disinfection.
4. To trap remaining particles, Cryptosporidium, E. coli and Giardia treat the pool with PRS in the following manner:

- a. Add PRS Stage 1 across the pool surface at a rate of 4 oz per 20,000 gallons of pool water. For best dispersion, add product in front of the pool return lines.
  
- b. Four hours after adding PRS Stage 1, add PRS Stage 2 across the pool surface at a rate of 4 oz per 20,000 gallons of pool water. For best dispersion, add product in front of the pool return lines.
  
- c. DO NOT add PRS Stage 1 and PRS Stage 2 at the same time.
  
- d. Monitor the filter pressure at regular intervals.
  
- e. Backwash the filter after proper a disinfection value is achieved.